Index

To Articles Published in The Nurse Practitioner in 1983

A

Albion, Johanna H., MS, MSN, CGNP: **Multiple Sclerosis: A Chronic-Care Focus,** Vol. 8, No. 4 (April 1983), p.29.

B

Baker, Clifford E., MA, RN, FNP, Crook, Jerry J., MD:
EKG of the Month, Vol. 8, No. 7 (July-August 1983), p. 12.
EKG of the Month, Vol. 8, No. 8 (September 1983), p. 68.
EKG of the Month, Vol. 8, No. 10 (November-December 1983), p. 78.

Barleben, Susan, RN, MSN, CFNP, Burns, Margaret, RN, MN, CFNP: **Preschooler and Adolescent: Repeating Themes,** Vol. 8, No. 8 (September 1983), p. 59.

Bartosch, Janet C., RNC, MS: **Oral Contraceptives: Selection and Management,** Vol. 8, No. 5 (May 1983), p. 56.

Briant, David. RN, MS, Brouder, Gerald, RN, PhD: **Identification and Treatment of Poision Ivy Dermatitis,** Vol. 8, No. 7 (July/August 1983), p. 13.

Brunt, J. Howard, RN: **Aspirin Therapy for Thromboembolic Disease,** Vol. 8, No. 2 (February 1983), p. 56.

C

Chalmers, Karen, Farrell, Patrica: **Nursing Interventions for Health Promotion,** Vol. 8, No. 10 (November-December 1983), p. 62.

Coslow, Florence, RN, BS, MA, CNP, Steinberg, Marion C., BS, MS: **Relaxation Techniques in Ambulatory Care Practice,** Vol. 8, No. 9 (October 1983), p. 26.

Coward, Doris D., BSN, RN: **The Other Herpesviruses: Epstein-Barr and Cytomegalovirus,** Vol. 8, No. 4 (April 1983), p. 13.

D

Doyle, Elaine, RNC, BSN, MEd, Meurer, Jeanne, RN, CNM, MSN: **Practicing Medicine Without a License,** Vol. 8, No. 6 (June 1983), p. 41.

Dunn, Ardys McNaughton: **Nurse Activism in Oregon Politics,** Vol. 8, No. 10 (November-December 1983), p. 54.

E

Edmunds, Marilyn, F[illegible], MS, ANP:
The Enthusiasm Plateau, Vol. 8, No. 2 (February 1983), p. 52.
The Nurse Preceptor Role, Vol. 8, No. 6 (June 1983), p. 49.
Models of Clinical Practice, Vol. 8, No. 9 (October 1983), p. 59.

F

Free, Teresa A., RN, PNP, MS: **The States' Challenge to Provide Maternal and Child Care,** Vol. 8, No. 5 (May 1983), p. 46.

Frenkel, Cheryl S., RN, MS, FNP, Pickett, Nancy G., RN, MS, FNP: **Political Awareness of Nurse Practitioners,** Vol. 8, No. 2 (February 1983), p. 9.

G

Gambert, Steven R., MD, FACP, Brensinger, John F. III RN, BSN, MS: **Assessing Thyroid Function in the Elderly,** Vol. 8, No. 7 (July/August 1983), p. 38.

Gaylord, Nan, RN, MS, CPNP, Carson, Sue, RN, BSN, CSN: **Assessing Recurrent Abdominal Pain in Children,** Vol. 8, No. 8 (September 1983), p. 19.

Gelfant, Barbara, RN, BSN, MSN: **Hypertension Protocol,** Vol. 8, No. 8 (September 1983), p. 25.

Giesbrecht, Velda, RNC, FNP: **Prevention of Taenia Saginata and Cysticerosis,** Vol. 8, No. 8 (September 1983), p. 56.

Gonzalez-Swafford, RN, MS, FNP, Gutierrez, Mary Grace, RN, MS, FNP: **Ethno-Medical Beliefs and Practices of Mexican-Americans,** Vol. 8, No. 10 (November-December 1983), p. 29.

Gorline, Lynn, RN, MS, FNC: **NPs in Tennessee: An Historical Overview,** Vol. 8, No. 10 (November-December 1983), p. 14.

Greene, Jane Cohen, RN, PNP, MN: **Prevention and Treatment of Sports Injuries,** Vol. 8, No. 10 (November-December 1983), p. 39.

H

Haggerty, Betty Jo, RN, MS, FNP: **Prevention and Differential of Scrotal Cancer,** Vol. 8, No. 10 (November-December 1983), p. 45.

Haggerty, Virginia C., RN, JD: **Doctrine of Delegated Medical Acts,** Vol. 8, No. 4 (April 1983), p. 9.

Hallman, Ellie C., RN, BN, Westlund Drexel, RN, BN: **Canadian Nurse Practitioners Battle Underutilization,** Vol. 8, No. 6 (June 1983), p. 45.

Harper, Doreen C., RN, C, PhD, Bilingsley, Molly C., RN, C, EdD: **Organizing for Power,** Vol. 8, No. 7 (July/August 1983), p. 24.

Heuther, Sue E., RN, PhD: **The Microcirculation,** Vol. 8, No. 7 (July/August 1983), p. 47.

Hilyard, Nancy, RN, MSN, CANP: **Solitary Thyroid Nodules in Adults,** Vol. 8, No. 2 (February 1983), p. 14.

Horn, Beverly, RN, PhD: **Cultural Beliefs and Teenage Pregnancy,** Vol. 8, No. 8 (September 1983), p. 35.

Hughes, Bobbie S., ARNP: **Role Evolution vs. Legislation,** Vol. 8, No. 3 (March 1983), p. 9.

J

Johnson, Jean, MSN, NP:
Medicare Cuts Hit Hospitals, Vol. 8, No. 1 (January 1983), p. 14.
The Saving of Social Security, Vol. 8, No. 5 (May 1983), p. 13.
Profile of a Nurse in the White House, Vol. 8, No. 8 (September 1983), p. 13.
Capitol Report, Vol. 8, No. 10 (November-December 1983), p. 17.

K

Karus, Celinda A., RN, BSN: **Tuberculosis: An Overview of Pathogenesis and Prevention,** Vol. 8, No. 2 (February 1983), p. 21.

Diet & Diabinese®
(chlorpropamide) Tablets 100 mg and 250 mg

A proven regimen for effective control of blood sugar.

BRIEF SUMMARY
DIABINESE® (chlorpropamide) Tablets

Contraindications: Diabinese is not indicated in patients having juvenile or growth-onset diabetes mellitus, severe or unstable "brittle" diabetes, and diabetes complicated by ketosis and acidosis, diabetic coma, major surgery, severe infection, or severe trauma.
Diabinese is contraindicated during pregnancy. Serious consideration should be given to the potential hazard of its use in women of childbearing age who may become pregnant.
Diabinese is contraindicated in patients with serious impairment of hepatic, renal, or thyroid function.
Precautions: Use chlorpropamide with caution with barbiturates, in patients with Addison's disease or in those ingesting: alcohol, antibacterial sulfonamides, thiazides, phenylbutazone, salicylates, probenecid, dicoumarol or MAO inhibitors. Adequate dietary intake should be assured in all patients using Diabinese.
Warnings: DIABINESE (CHLORPROPAMIDE) SHOULD NOT BE USED IN JUVENILE DIABETES OR IN DIABETES COMPLICATED BY ACIDOSIS, COMA, SEVERE INFECTION, MAJOR SURGICAL PROCEDURES, SEVERE TRAUMA, SEVERE DIARRHEA, NAUSEA AND VOMITING, ETC. HERE, INSULIN IS INDISPENSABLE.
HYPOGLYCEMIA, IF IT OCCURS, MAY BE PROLONGED. (SEE ADVERSE REACTIONS.) IN INSTANCES OF CONCOMITANT USE WITH INSULIN, PATIENTS SHOULD BE CAREFULLY MONITORED.
Adverse Reactions: Usually dose-related and generally respond to reduction or withdrawal of therapy. Generally transient and not of a serious nature and include anorexia, nausea, vomiting and gastrointestinal intolerance; weakness and paresthesias.
Certain untoward reactions associated with idiosyncrasy or hypersensitivity have occasionally occurred, including jaundice, skin eruptions rarely progressing to erythema multiforme and exfoliative dermatitis, and probably depression of formed elements of the blood. They occur characteristically during the first six weeks of therapy. With a few exceptions, these manifestations have been mild and readily reversible on the withdrawal of the drug. The more severe manifestations may require other therapeutic measures, including corticosteroid therapy. Diabinese should be discontinued promptly when the development of sensitivity is suspected.
Jaundice has been reported, and is usually promptly reversible on discontinuance of therapy. THE OCCURRENCE OF PROGRESSIVE ALKALINE PHOSPHATASE ELEVATION SHOULD SUGGEST THE POSSIBILITY OF INCIPIENT JAUNDICE AND CONSTITUTES AN INDICATION FOR WITHDRAWAL OF THE DRUG.
Leukopenia, thrombocytopenia and mild anemia, which occur occasionally, are generally benign and revert to normal, following cessation of the drug.
Cases of aplastic anemia and agranulocytosis, generally similar to blood dyscrasias associated with other sulfonylureas, have been reported.
BECAUSE OF THE PROLONGED HYPOGLYCEMIC ACTION OF DIABINESE, PATIENTS WHO BECOME HYPOGLYCEMIC DURING THERAPY WITH THIS DRUG REQUIRE CLOSE SUPERVISION FOR A MINIMUM PERIOD OF 3 TO 5 DAYS, during which time frequent feedings or glucose administration are essential. The anorectic patient or the profoundly hypoglycemic patient should be hospitalized.
Rare cases of phototoxic reactions have been reported. Edema associated with hyponatremia has been infrequently reported. It is usually readily reversible when medication is discontinued.
Dosage: The total daily dosage is generally taken at a single time each morning with breakfast. Occasionally, cases of gastrointestinal intolerance may be relieved by dividing the daily dosage. A LOADING OR PRIMING DOSE IS NOT NECESSARY AND SHOULD NOT BE USED. The mild to moderately severe, middle-aged, stable diabetic should be started on 250 mg daily. Because the geriatric diabetic patient appears to be more sensitive to the hypoglycemic effect of sulfonylurea drugs, older patients should be started on smaller amounts of Diabinese, in the range of 100 to 125 mg daily.
After five to seven days following initiation of therapy, dosage may be adjusted upward or downward in increments of 50 to 125 mg at intervals of three to five days. PATIENTS WHO DO NOT RESPOND COMPLETELY TO 500 MG DAILY WILL USUALLY NOT RESPOND TO HIGHER DOSES. Maintenance doses above 750 mg daily should be avoided.
Supply: 100 mg and 250 mg, blue, 'D'-shaped, scored tablets.

More detailed professional information available on request.

Pfizer LABORATORIES DIVISION
PFIZER INC.

Leaders in Oral Diabetic Therapy

Keller, Elizabeth K., RN, BSN: **Patellar Malalignment Syndrome in Runners,** Vol. 8, No. 6 (June 1983), p. 27.

Kilcoyne, Raphael F., MD, Plumley, Thomas F., MD, Marglin, Stephen I., MD: **Infections of Bones and Joints,** Vol. 8, No. 3 (March 1983), p. 63.

L

Lamberton, Martha M., RN, MSN: **Health and Illness: A Co-Existence Hypothesis,** Vol. 8, No. 2 (February 1983), p. 47.

Lauver, Diane, FNPC: **Childhood Enuresis,** Vol. 8, No. 2 (February 1983), p. 30.

Leff, Susan D., RN, PHN: **The Question of a Standard of Care,** Vol. 8, No. 7 (July/August 1983), p. 34.

Libbus, M. Kay, MSPH, RN, C, FNP: **Enterobiasis,** Vol. 8, No. 8 (September 1983), p. 17.

Loucks, Arlene, RNC, MSN: **Pelvic Inflammatory Disease: A Review of Therapy,** Vol. 8, No. 9 (October 1983), p. 13.

Luce, Margaret, RN, MSN, Sande, Diane, RN, MS: **Oral Health in Children: Prevention of Dental Caries,** Vol. 8, No. 3 (March 1983),

M

Macek, Loretta, RN, BSN: **Acute Rheumatic Fever,** Vol. 8, No. 5 (May 1983), p. 17.

Marglin, Stephen I., MD: **Atelectasis,** Vol.8, No. 1 (January 1983), p. 13.

Maxwell, Betty McCloud, RN, MSN: **The Special Olympics,** Vol. 8, No. 10 (November-December 1983), p. 65.

McCance, Kathryn L., RN, PhD: **Lipoproteins and Heart Disease,** Vol. 8, No. 5 (May 1983), p. 68.

McCarthy, Patricia A., RN, MS, CFNP: **Fetal Alcohol Syndrome and Other Alcohol-Related Birth Defects,** Vol. 8, No. 1 (January 1983), p. 33.

Moniz, Donna M., CRN, MN, JD, Tarutis, Gerald R., JD: **Making Law, Not Breaking Law,** Vol. 8, No. 9 (October 1983), p. 66.

N

Nuccio, Maureen A., Colleen Sparks: **Intestinal Parasites: Diagnostic and Treatment Guidelines,** Vol. 8, No. 1 (January 1983), p. 47.

P

Parker, Susan O., RN, C, MSN: **A Conceptual Model for Outcome Assessment,** Vol. 8, No. 1, (January 1983), p. 41.

Pearson, Linda, RN, MSN, CFNP: **Is a Nurse Ever Not a Nurse?,** Vol. 8, No. 6 (June 1983), p. 8.

Pickwell, Sheila, MN, CFNP: **Protocol: Health Screening for Indochinese Refugees,** Vol. 8, No. 4 (April 1983), p. 20.

Plein, Joy B., RPh, PhD: **The Penicillins,** Vol. 8, No. 1 (January 1983), p. 51.

Poole, Anita L., MSN, PNP, Poole, Steven R., MD: **The Nurse Practitioner's Role in Adolescent Health Care,** Vol. 8, No. 9 (October 1983), p. 43.

Power, Robin, RN, BSN: **Hereditary Polymorphic Light Eruption,** Vol. 8, No. 6 (June 1983), p. 19.

Q

Quinn, Nancy, BN, MS, AHNP: **Health Hazards of Operating Video Display Terminals,** Vol. 8, No. 9 (October 1983), p. 49.

R

Rapson, Mary F., RN, PhD, CANP: **Strategies for Coping with Role Stress,** Vol. 8, No. 5 (May 1983), p. 75.

NALDECON® antihistamine/decongestant
Tablets, Syrup, Pediatric Drops and Pediatric Syrup

FOR ORAL USE ONLY

For complete information consult Official Package Circular.

CONTRAINDICATIONS: Patients with severe hypertension, severe coronary artery disease, patients on MAO inhibitor therapy; patients with narrow angle glaucoma, urinary retention, peptic ulcer and during an asthmatic attack. Also contraindiated in patients with hypersensitivity or idiosyncrasy to sympathomimetic amines or antihistamines.

WARNINGS: Sympathomimetic amines should be used judiciously and sparingly in patients with hypertension, diabetes mellitus, ischemic heart disease, increased intraocular pressure, hyperthyroidism or prostatic hypertrophy. See, however, CONTRAINDICATIONS. Sympathomimetics may produce central nervous system stimulation with convulsions or cardiovascular collapse with accompanying hypotension.

Antihistamines may impair mental and physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery, and may impair mental alertness in children. Chlorpheniramine and phenyltoloxamine have an atropine-like action and should be used with caution in patients with increased intraocular pressure, cardiovascular disease, hypertension or in patients with a history of bronchial asthma. See, however, CONTRAINDICATIONS.

Do not exceed recommended dosage.

PRECAUTIONS: Patients with diabetes, hypertension, cardiovascular disease and hyperreactivity to ephedrine. The antihistaminics may cause drowsiness and ambulatory patients who operate machinery or motor vehicles should be cautioned accordingly.

DRUG INTERACTIONS: MAO inhibitors and beta adrenergic blockers increase the effect of sympathomimetics. Sympathomimetics may reduce the antihypertensive effects of methyldopa, mecamylamine, reserpine and veratrum alkaloids. Concomitant use of antihistamines with alcohol, tricyclic antidepressants, barbiturates and other CNS depressants may have an addictive effect.

PREGNANCY CATEGORY C: Animal reproduction studies have not been conducted with Naldecon. It is also not known whether Naldecon can cause fetal harm when administered to a pregnant woman or can effect reproductive capacity. Naldecon should be given to a pregnant woman only if clearly needed.

NURSING MOTHERS: Caution should be exercised when this drug is given to nursing mothers due to the higher than usual risk of the sympathomimetic amines in infants.

ADVERSE REACTIONS: Hyperreactive individuals may display ephedrine-like reactions such as tachycardia, palpitations, headache, dizziness or nausea. Patients sensitive to antihistamines may experience mild sedation. Sympathomimetics have been associated with certain untoward reactions including restlessness, tremor, weakness, pallor, respiratory difficulty, dysuria, insomnia, hallucinations, convulsions, CNS depression, arrhythmias and cardiovascular collapse with hypotension. Possible side effects of antihistamines are drowsiness, restlessness, dizziness, weakness, dry mouth, anorexia, nausea, vomiting, headache, nervousness, blurring of vision, polyuria, heartburn, dysuria and, very rarely, dermatitis.

HOW SUPPLIED: NDC 0015-5600—Naldecon Tablets; **NDC** 0015-5601—Naldecon Syrup; **NDC** 0015-5615—Naldecon Pediatric Drops; **NDC** 0015-5616—Naldecon Pediatric Syrup

For information on package sizes available, refer to the current price schedule.

NALDECON—(11)—7/3/80

BRISTOL™

©1983 BRISTOL LABORATORIES
Division of Bristol-Myers Company
Syracuse, New York 13221-4755

BL-1812 9/83

Reid, Jacquelyn D., RN, C, EdD: **Effects of Selected OTC Medications on the Unborn and Newborn,** Vol. 8, No. 8 (September 1983), p. 43.

Roberts, Leslie J., RN, MHSC, Ross, Eleanor, RN, BSN, MSN: **Chronic Airflow Limitation,** Vol. 8, No. 10 (November-December 1983), p. 21.

Roos, Philip D., PhD, Crooker, Margaret, RN, MSN, CPNP: **Variables Affecting Nurse Practitioner Salaries,** Vol. 8, No. 5 (May 1983), p. 36.

Runtz, Susan E., RN, MEd, MSN, CANP, Urtel, Janice G., RN, MA, MSN, CANP: **Evaluating Your Practice Via a Nursing Model,** Vol. 8, No. 3 (March 1983), p. 30.

S

Schumann, Delores: **Diabetes Mellitus and Exercise: Application of Research to Activity Planning,** Vol. 8, No. 3 (March 1983), p. 13.

Sears, Helen, RN, BSN, Morken, Jean, RN, BSN: **Kaposi's Sarcoma: A Deadly New Syndrome,** Vol. 8, No. 1 (January 1983), p. 25.

Shortridge, Lillie M., RN, EdD, FNC, FAAN, McLain, Barbara R., RN, MS, NP: **Levels of Intervention for a Coexistence Model,** Vol. 8, No. 3 (March 1983), p. 74.

Stillwell, Susan B., RN, MSN: **The Edematous Client: Causes, Physical Assessment and Treatment,** Vol. 8, No. 3 (March 1983), p. 21.

Stockdale-Woolley, Rebecca, RN, MSN: **Sexual Dysfunction and COPD: Problems and Management,** Vol. 8, No. 2 (February 1983), p. 16.

Sylvester, Carol, RN, MS, ANP-C, Vish, Denise, RN, MS, ANP-C, Edmunds, Marilyn W., RN, MS, ANP-C: **Site Assessment for NPs in the Geriatric Setting,** Vol. 8, No. 7 (July/August 1983), p. 54.

T

Talley, Sandra, RN, MN, CS:
- **Bridging Psychiatric and Primary Care,** Vol. 8, No. 3 (March 1983), p. 70.
- **Psychiatric vs. Psychosocial Concepts,** Vol. 8, No. 5 (May 1983), p. 83.
- **Intervention Styles for Psychiatric and Primary Care,** Vol. 8, No. 6 (June 1983), p. 56.
- **Psychiatric Consultation and Collaboration,** Vol. 8, No. 9 (October 1983), p. 55.

Thomas, Courtney, RN, MSN, PNP, CFNP:
- **Scope of Practice Issues,** Vol. 8, No. 1 (January 1983), p. 21.
- **–State Report,** Vol. 8, No. 5 (May 1983), p. 12.
- **–State Legislative Activities Reflect NP Efforts,** Vol. 8, No. 7 (July/August 1983), p. 9.
- **–State Report,** Vol. 8, No. 9 (October 1983), p. 12.

Trandel-Korenchuk, Darlene, RN, MSN, FNP, Trandel-Korenchuk, Keith, JD, MPH:
- **Part 1, Nursing Advocacy of Patients' Rights: Myth or Reality?** Vol. 8, No. 3 (March 1983), p. 53.
- **Part 2, Nursing Advocacy of Patients' Rights: Myth or Reality?,** Vol. 8, No. 4 (April 1983), p. 37.

W

Webster-Stratton, Carolyn, PhD: **Intervention Approaches to Conduct Disorders in Young Children,** Vol. 8, No. 5 (May 1983), p. 23.

Williams, Ann, FNP: **Hepatitis B Virus and Vaccine,** Vol. 8, No. 9 (October 1983), p. 30.

Wolbert, Jerry, RN, MS, FNP: **Sexually Transmitted Diseases in Homosexual Men,** Vol. 8, No. 9 (October 1983), p. 35.

Y

Young, Daryl, RN, MS, FNP: **Emergency Intervention in Traumatic Amputation,** Vol. 8, No. 10 (November-December 1983), p. 57.